

ICESB 2011: 25-26 November 2011, Maldives

Effect of the Extract of *Fructus Tribuli* on Growth of *Lactobacillus Acidophilus*

Shu Guowei^{a,b,*}, Ji Liyuan^a, Qin Tao^b, Chen He^a, Ma Qi^b

^aCollege of Life Science & Engineering, Shaanxi University of Science & Technology, Xi'an, 710021, China

^bEnzyme Engineering Institute, Shaanxi Academy of Sciences, Xi'an, 710021, China

Abstract

Effect of the extract of *Fructus tribuli* (EFT) on growth of *Lactobacillus acidophilus* LA04, LA05, LA06 and LA09 was studied by measuring optical density at 600nm (OD600) and pH using MRS media as the control. The addition of EFT (v/v) was 1%, 2%, 3%, 4% and 5%. Results were as follows: addition of EFT could promote the growth of *Lactobacillus acidophilus* LA09, The optimum concentration of EFT in MRS media was 1% at incubation 24h, but addition of EFT on growth of *Lactobacillus acidophilus* LA04, LA05 and LA06 has no significant effect.

© 2011 Published by Elsevier Ltd. Selection and/or peer-review under responsibility of the Asia-Pacific Chemical, Biological & Environmental Engineering Society (APCBEEES) Open access under [CC BY-NC-ND license](#).

Keywords: *Lactobacillus acidophilus*; probiotics; *Fructus tribuli*;

1. Introduction

The probiotics are “Live microorganisms, which when administered in adequate amounts confer a health benefit on the host”. The probiotics that have been traditionally most often used are the lactic bacteria of the genera *Lactobacillus*, *Streptococcus*, the *Bifidobacterium* plus *Saccharomyces boulardii*, which have been introduced in food products and drugs due to their growing evidence of health benefits [1-2]. Strains of *Lactobacillus acidophilus* and *Lactobacillus rhamnosus* strain GG probably have the longest history of application as probiotics. Clinical studies have also supported the role of probiotics in treating pouchitis, antibiotic-associated diarrhea (AAD) and travelers' diarrhea, Crohn's disease, ulcerative colitis (UC), irritable bowel syndrome (IBS), *Helicobacter pylori* infections, lactose intolerance, immunomodulation, lowering of blood cholesterol [3-8].

* Corresponding author. Tel.: +0086-029-86168589; fax: +0086-029-86168293.

E-mail address: shuguowei@gmail.com.

Tribulus terrestris is an annual herb that grows all over the world [9]. The fruit, stems and leaves of the plant are used in the folk medicine in China against various diseases, including impotence tonic, cardiovascular diseases, anti-hypertensive urinary anti-infective, anti-inflammatory and immunosuppressive activities [10-11]. Fructus tribuli is the fruit of *Tribulus terrestris* and allowed to use in functional food by the Ministry of Health of China. Furostanol and spirostanol saponins, polysaccharides, flavonoid glycosides, alkaloids, alkaloids and some amides have been reported to occur in Fructus Tribuli [12-15].

In previous research, we found that the hot water extract from Fructus tribuli can promote the growth of *L. acidophilus* WS [16]. The purpose of the present work was to study effect of EFT on growth of *Lactobacillus acidophilus* LA04, LA05, LA06 and LA09.

2. Materials and methods

2.1. Microorganism

Lactobacillus acidophilus LA04, LA05, LA06 and LA09 were obtained from College of Life Science & Engineering, Shaanxi University of Science & Technology. All lactobacilli strains were grown three successive times in MRS medium (Hopebio, Qingdao, China) before use.

2.2. Preparation of the extract from Fructus tribuli

Dried *Fructus tribuli* were purchased from a local herb market (Xi'an, China), *Fructus tribuli* was ground in a high speed disintegrator (Model SF-2000, Shanghai, China) to obtain a fine powder, then was extracted distilled water (liquid: solid ratio (ml/g) 14:1, While the temperature of the water bath was kept steady, The extraction in a 1.0 l stainless steel boiler in the water bath was stirred with an electric mixing paddle for 1.5h, then obtained the extract of Fructus tribuli (EFT) by filtration.

2.3. Culture conditions

EFT was added to MRS media in anaerobic tube at 1.0%, 2.0%, 3.0%, 4.0% and 5.0% (v/v), respectively. MRS media without EFT was included in this experiment as a control.

All samples were autoclaved at 118°C for 15 min. Active culture was added to each anaerobic tube after cooling to room temperature and incubated at 37°C, then determined biomass and pH at incubation 12h, 18h and 24h. The experiment was completed in triplicate and means were calculated for each treatment group at each data collection point.

2.4. Biomass and pH measurement

The biomass was determined by optical density at 600nm (OD₆₀₀) through a spectrophotometer (SP-756PC, Shanghai Spectrum Instruments Co., Ltd., Shanghai, China). The pH of culture was measured through a pH-meter (PHS-3C Shanghai Precision Scientific Instrument Co., Ltd, Shanghai, China).

3. Results and discussion

3.1. Effect of EFT on growth of *L. acidophilus* LA04

Effect of different concentrations of EFT on the growth of *Lactobacillus acidophilus* LA04 showed in Figure 1. With the increasing concentration of EFT, OD value increased gradually and reached maximum 1.605, 1.788 and 1.772 at 1%, 4% and 2% EFT, then decreased to 1.542, 1.728 and 1.720 at 5% EFT at incubation 12h, 18h and 24h, respectively. The OD value of the control was 1.574, 1.726 and 1.736, respectively; the pH of different culture mediums has no significant difference ranging from 4.06 to 4.12(incubation time 12h), 3.79-3.84(incubation time 18h) and 3.70-3.76(incubation time 24h), which indicated that EFT on growth of *Lactobacillus acidophilus* LA04 has no significant role.




Fig1. Effect of EFT on growth of *Lactobacillus acidophilus* LA04, A, B and C mean incubation 12h, 18h and 24h, respectively

3.2. Effect of EFT on growth of *L. acidophilus* LA05

Effect of different concentrations of EFT on the growth of *Lactobacillus acidophilus* LA05 showed in Figure 2. With the increasing concentration of EFT, OD value increased gradually and reached maximum 1.629, 1.797 and 1.783 at 1% EFT, then decreased to 1.493, 1.575 and 1.539 at 5% EFT at incubation 12h, 18h and 24h, respectively. The OD value of the control was 1.580, 1.746 and 1.734, respectively; the pH of different culture mediums in contrast with the OD₆₀₀ has no significant difference ranging from 4.10 to 4.28(incubation time 12h), 3.77-4.12(incubation time 18h) and 3.70-4.04(incubation time 24h), which indicated that EFT on growth of *Lactobacillus acidophilus* LA05 has no obvious effect.

3.3. Effect of EFT on growth of *L. acidophilus* LA06

Effect of different concentrations of EFT on the growth of *Lactobacillus acidophilus* LA06 showed in Figure 3. With the increasing concentration of EFT, OD value increased gradually and reached maximum 1.627 and 1.642 at 2% and 1% EFT, then decreased to 1.567 and 1.61 at 5% EFT at incubation 12h and 18h, respectively. The OD value increased gradually from 1.603 to 1.621 at 5% EFT at incubation 24h, The OD₆₀₀ of the control was 1.567, 1.616 and 1.578, respectively; the pH of different culture mediums in contrast with the OD₆₀₀ has no significant difference ranging from 4.10 to 4.28(incubation time 12h), 3.77-4.12(incubation time 18h) and 3.70-4.04(incubation time 24h), which indicated that EFT on growth of *Lactobacillus acidophilus* LA06 has no significant role.




Fig2. Effect of EFT on growth of *Lactobacillus acidophilus* LA05, A, B and C mean incubation 12h, 18h and 24h, respectively





Fig3. Effect of EFT on growth of *Lactobacillus acidophilus* LA06, A, B and C mean incubation 12h, 18h and 24h, respectively



3.4. Effect of EFT on growth of *L. acidophilus* LA09

Effect of different concentrations of EFT on the growth of *Lactobacillus acidophilus* LA09 showed in Figure 4. With the increasing concentration of EFT at incubation 12h, OD value increased gradually from 0.655 at 0% EFT to 1.042 at 3% EFT, then decreased to 1.034 at 5% EFT. At incubation 18h, The OD value increased gradually from 0.914 at 0% EFT to 1.181 at 1% EFT, then decreased to 1.115 at 5% EFT. At incubation 24h, The OD value increased gradually from 1.027 at 0% EFT to 1.265 at 1% EFT, then decreased to 1.141 at 5% EFT. the pH of different culture mediums in contrast with the OD₆₀₀ has significant difference ranging from 5.01 to 5.17 (incubation time 12h), 4.79-5.09 (incubation time 18h and 24h), which indicated that EFT on growth of *Lactobacillus acidophilus* LA09 has an significant promotion, and the optimum concentration of EFT in MRS media was 1%.




Fig4. Effect of EFT on growth of *Lactobacillus acidophilus* LA09, A, B and C mean incubation 12h, 18h and 24h, respectively

4. Conclusions

Addition of EFT in MRS media has the significant promotion on growth of *Lactobacillus acidophilus* LA09, but has no obvious effect to *Lactobacillus acidophilus* LA04, LA05 and LA06. The optimum concentration of EFT in MRS media was 1% for *Lactobacillus acidophilus* LA09.

Acknowledgements

The project was supported by the key project of Shaanxi Academy of Sciences (No.2010K-03), the Natural Science Foundation of Shaanxi Province (No.2010JM2013), key project of Agriculture Department of Shaanxi Province (No.2009-45), Science and Technology Bureau of Weiyang District (No.201002).

References

- [1] Nancy Toedter Williams. Probiotics. *Am J Health-Syst Pharm*. 2010;**67**:449-59.
- [2] Kalliomaki, M., Salminen, S., Poussa, T., Arvilommi, H. Probiotics and prevention of atopic disease: 4-year follow-up of a randomised placebo-controlled trial. *Lancet*, 2003;**361**:1869-71.
- [3] Senok AC, Ismaeel AY, Botta GA. Probiotics: facts and myths. *Clin Microbiol Infect* 2005;**11**:958-66
- [4] Doron S, Gorbach SL. Probiotics: their role in the treatment and prevention of disease. *Expert Rev Anti Infect Ther* 2006;**4**: 261-75.
- [5] Wald A, Rakel D. Behavioral and complementary approaches for the treatment of irritable bowel syndrome. *Nutr Clin Pract* 2008;**123**:284-92.
- [6] Surawicz CM. Role of probiotics in antibiotic-associated diarrhea, *Clostridium difficile*-associated diarrhea, and recurrent *Clostridium difficile*-associated diarrhea. *J Clin Gastroenterol* 2008;**42**:S64-70.
- [7] MacIntyre A, Cymet TC. Probiotics: the benefits of bacterial cultures. *Compr Ther* 2005;**31**:181-5.
- [8] Farnworth ER. The evidence to support health claims for probiotics. *J Nutr* 2008;**138**:1250-4.
- [9] Combarieu E.D, Fuzzati N, Lovati M, calli E.M. Furostanol saponins from *Tribulus terrestris*. *Fitoterapia* 2003;**74**: 583–91.
- [10] Ali M. Sharifi, Radbod Darabi, Nasrin Akbarloo. Study of antihypertensive mechanism of *Tribulus terrestris* in 2K1C hypertensive rats: Role of tissue ACE activity. *Life Sciences* 2003;**73**:2963–71.
- [11] Joshia V.S., Parekha B.B., Joshia M.J.& Vaidya A.B. Herbal extracts of *Tribulus terrestris* and *Bergenia ligulata* inhibit growth of calcium oxalate monohydrate crystals in vitro. *Journal of Crystal Growth* 2005;**275**:1403–8.
- [12] Fang S., Hao C., Liu Z., Song F., Liu S. Application of electrospray ionization mass spectrometry combined with sequential tandem mass spectrometry techniques for the profiling of steroidal saponin mixture extracted from *Tribulus terrestris*. *Planta Med* 1999;**65**:68–73.
- [13] Kostova I, Dinchev D, Rentsch GH, Dimitrov V, Ivanova A. Two New Sulfated Furostanol Saponins from *Tribulus terrestris*. *Z Naturforsch* 2002;**57**:33-8.
- [14] Kadota S. Tribulusamide A and B, New hepatoprotective lignanamides from the fruits of *Tribulus terrestris*. Indications of cytoprotective activity in murine hepatocyte culture. *Planta Med* 1998;**64**: 628–31.
- [15] Wu T.S., Shi L.S., Kuo S.C. Alkaloids and other constituents from *Tribulus terrestris*. *Phytochemistry* 1999;**50**:1411-5.
- [16] Shu.G.W. Lu.J.L. Chen H. Research on Reciprocity between Probiotics and *Tribulus Terrestris*. *Science and Technology of Food Industry* 2007;**31**:103-5.